

On the algebraic solutions of the sixth Painlevé equation related to second order Picard-Fuchs equations

Bassem BEN HAMED[†] Lubomir GAVRILOV[‡]

February 1, 2008

[†] Faculté des Sciences de Sfax, Département de Mathématiques
BP 802, route Soukra km 4 Sfax 3018, Tunisie

[‡] Laboratoire Emile Picard, CNRS UMR 5580, Université Paul Sabatier
118, route de Narbonne, 31062 Toulouse Cedex 4, France

Abstract

We describe two algebraic solutions of the sixth Painlevé equation which are related to (isomonodromic) deformations of Picard-Fuchs equations of order two.

1 Statement of the result

In this note we describe two algebraic solutions of the following Painlevé VI (\mathcal{P}_{VI}) equation

$$\begin{aligned} \frac{d^2\lambda}{dt^2} = & \frac{1}{2}\left(\frac{1}{\lambda} + \frac{1}{\lambda-1} + \frac{1}{\lambda-t}\right)\left(\frac{d\lambda}{dt}\right)^2 - \left(\frac{1}{t} + \frac{1}{t-1} + \frac{1}{\lambda-t}\right)\frac{d\lambda}{dt} \\ & + \frac{\lambda(\lambda-1)(\lambda-t)}{t^2(t^2-1)}\left[\alpha + \beta\frac{t}{\lambda^2} + \gamma\frac{t-1}{(\lambda-1)^2} + \delta\frac{t(t-1)}{(\lambda-t)^2}\right]. \end{aligned}$$

related to deformations of Picard-Fuchs equations of special type. Recall that the \mathcal{P}_{VI} equation governs the isomonodromic deformations of the second

order Fuchsian equations

$$x'' + p_1(s)x' + p_2(s)x = 0, \quad ' = \frac{d}{ds} \quad (1)$$

with 5 singular points, one of which is apparent [7]. Suppose that the solution of the Fuchsian equation (1) is given by an Abelian integral

$$x(s) = \int_{\gamma_s} \omega$$

where ω is a rational one-form on \mathbb{C}^2 , $\Gamma_s \subset \mathbb{C}^2$ is a family of algebraic curves depending rationally on s , and $\gamma_s \subset \Gamma_s$ is a continuous family of closed loops. Then the equation (1) is said to be of Picard-Fuchs type and its monodromy group is conjugated to a subgroup of $\mathbf{GL}_2(\overline{\mathbb{Q}})$ (generically $\mathbf{GL}_2(\mathbb{Z})$). For this reason any continuous deformation

$$a \rightarrow \Gamma_{s,a}$$

of the family Γ_s induces an isomonodromic deformation of (1). If in addition $\Gamma_{s,a}$ depends algebraically in a , the coefficients of (1) are also algebraic functions in a , and hence they provide an algebraic solution of \mathcal{P}_{VI} .

From now on we denote

$$(\alpha_0, \alpha_1, \alpha_2, \alpha_3) = (\alpha, -\beta, \gamma, \frac{1}{2} - \delta).$$

Our main result is the following

Theorem 1 *The pencil of $\mathcal{P}_{VI}(\alpha)$ equations*

$$(\alpha_0, \alpha_1, \alpha_2, \alpha_3) = (\frac{1}{8}, \frac{s}{8}, \frac{s}{8}, \frac{s}{8}), s \in \mathbb{C} \quad (2)$$

has a common algebraic solution parameterized as

$$\lambda = \frac{a^2(2-a)}{a^2-a+1}, t = \frac{a^3(2-a)}{2a-1}, a \in \mathbb{C}. \quad (3)$$

The $\mathcal{P}_{VI}(\alpha)$ equation with

$$\alpha = (\frac{1}{8}, \frac{1}{2}, 0, 0)$$

has an algebraic solution parameterized as

$$\lambda = \frac{a(a-2)(2a^2+a+2)}{a^2-7a+1}, t = \frac{a^3(2-a)}{2a-1}, a \in \mathbb{C} \quad (4)$$

The meaning of these solutions is the following. Consider the family of elliptic curves

$$\Gamma_s = \{(\xi, \eta) \in \mathbb{C}^2 : \eta^2 + \frac{3}{2a-1}\xi^4 - \frac{4(a+1)}{2a-1}\xi^3 + \frac{6a}{2a-1}\xi^2 = s\} \quad (5)$$

and let $\gamma(s) \in H_1(\Gamma_s, \mathbb{Z})$ be a family of cycles depending continuously on $s \in \mathbb{C}$. The Abelian integral of first kind

$$\int_{\gamma(s)} \frac{d\xi}{\eta}$$

satisfies a Picard-Fuchs equation of second order depending on a parameter a , defining an isomonodromy deformation of the equation. This deformation corresponds then to an algebraic solution of $\mathcal{P}_{VI}(\alpha)$ given by (3). In a similar way, the Abelian integral of second kind

$$\int_{\gamma(s)} \frac{(3\xi^2 - 2(a+1))\xi d\xi}{\eta}$$

satisfies a Picard-Fuchs equation of second order. The isomonodromy deformation of this equation with respect to a is described by the solution (4) of \mathcal{P}_{VI} equation.

Algebraic solutions of \mathcal{P}_{VI} were found by many authors, e.g. Hitchin[6], Manin[8], Dubrovin-Mazzocco[4], Boalch[3]. Dubrovin and Mazzocco classified all algebraic solutions of the \mathcal{P}_{VI} equation corresponding to

$$(\alpha_0, \alpha_1, \alpha_2, \alpha_3) = \left(\frac{1}{2}(2\mu - 1)^2, 0, 0, 0\right), \mu \in \mathbb{R}.$$

It turns out that these solutions, up to symmetries, are in a one-to-one correspondence with the regular polyhedra in the three dimensional space. Our solution (3) with

$$(\alpha_0, \alpha_1, \alpha_2, \alpha_3) = (1/8, 0, 0, 0)$$

corresponds then to the tetrahedron solution of Dubrovin-Mazzocco ($\mu = +1/4$). It is identified to their solution (A_3) via the Okamoto type transformation (1.24),(1.25), see [4]. It is remarkable that the same solution, but for

$$(\alpha_0, \alpha_1, \alpha_2, \alpha_3) = (1/8, 1/8, 1/8, 1/8)$$

was also found by Hitchin[6]. This shows that (3) is a common solution to the family (2) of \mathcal{P}_{VI} equations. It is clear that for transcendental values of

s in (2) the corresponding isomonodromic family of Fuchs equations (1) can not be of Picard-Fuchs type.

The paper is organized as follows. In the next section we recall briefly, following [7], the relationship between \mathcal{P}_{VI} and the isomonodromic deformations of Fuchs equations. In section 3 we deduce the relevant Picard-Fuchs equation and establish the main result.

The present text is an abridged version of [2].

2 The Garnier system and the \mathcal{P}_{VI} equation.

Consider a Fuchsian differential equation

$$x'' + p_1(s)x' + p_2(s)x = 0, \quad ' = \frac{d}{ds} \quad (1)$$

with *five singular points, exactly one of which is apparent*. After a bi-rational change of the independent variable s and a linear change of the dependent variable x (involving s) we may suppose that the singular points are $0, 1, t, \lambda, \infty$, where the singularity λ is apparent and the corresponding Riemann scheme is

$$\begin{pmatrix} 0 & 1 & t & \lambda & \infty \\ 0 & 0 & 0 & 0 & \alpha \\ \theta_1 & \theta_2 & \theta_3 & k & \alpha + \theta_\infty \end{pmatrix}, n \in \mathbb{N}, 2\alpha + \sum_i \theta_i + n = 3.$$

In what follows we shall always suppose that $n = 2$ (which is satisfied generically)).

The coefficients p_1, p_2 are easily computed to be

$$p_1(s) = \frac{1 - \theta_1}{s} + \frac{1 - \theta_2}{s - 1} + \frac{1 - \theta_3}{s - t} - \frac{1}{t - \lambda}$$

$$p_2(s) = \frac{k}{s(s - 1)} - \frac{t(t - 1)K}{s(s - 1)(s - t)} + \frac{\lambda(\lambda - 1)\mu}{s(s - 1)(s - \lambda)}$$

where μ is a constant

$$k = \frac{1}{4} \left\{ \left(\sum_{i=1}^3 \theta_i - 1 \right)^2 - \theta_\infty^2 \right\}.$$

The compatibility condition for the singular point λ to be apparent reads

$$K = K(\lambda, \mu, t) = \frac{1}{t(t-1)}[\lambda(\lambda-1)(\lambda-t)\mu^2 - \{\theta_2(\lambda-1)(\lambda-t) + \theta_3\lambda(\lambda-t) + (\theta_1-1)\lambda(\lambda-1)\}\mu + k\lambda].$$

From the discussion above it is seen that the Fuchs equation (1) depends on the parameters $\theta_0, \theta_1, \theta_t, \theta_\infty, \lambda, \mu, t$. Let us denote this equation by $E_\theta(\lambda, \mu, t)$.

Theorem 2 $\lambda(t), \mu(t)$ is a solution of the Garnier system

$$\begin{aligned} \frac{d\lambda}{dt} &= \frac{\partial K}{\partial \mu} \\ \frac{d\mu}{dt} &= -\frac{\partial K}{\partial \lambda}. \end{aligned}$$

if and only if the induced deformation of $E_\theta(\lambda, \mu, t)$ is isomonodromic.

It is straightforward to check that the sixth Painlevé system $\mathcal{P}_{VI}(\alpha)$ with parameters

$$\alpha = (\frac{1}{2}\theta_\infty^2, \frac{1}{2}\theta_0^2, \frac{1}{2}\theta_1^2, \frac{1}{2}\theta_t^2) \quad (6)$$

is equivalent to the Garnier system. We get therefore the following

Corollary. If

$$(t, \lambda, \mu) \rightarrow (t, \lambda(t), \mu(t))$$

is an isomonodromic deformation of $E_\theta(\lambda, \mu, t)$, then $\lambda(t)$ is a solution of $\mathcal{P}_{VI}(\alpha)$ equations with parameters given by (6).

3 Picard-Fuchs equations

In this section we restrict our attention to the deformation

$$f_a(\xi, \eta) = \eta^2 + \frac{3}{2a-1}\xi^4 - \frac{4(a+1)}{2a-1}\xi^3 + \frac{6a}{2a-1}\xi^2, a \in \mathbb{C}$$

of the singularity $\eta^2 + \xi^4$ of type A_3 , see [1]. The critical values of $f_a(\xi, \eta)$ are

$$0, 1, t = \frac{a^3(2-a)}{2a-1}.$$

Consider the locally trivial smooth fibration

$$f^{-1}(\mathbb{C} \setminus \{0, 1, t\}) \rightarrow \mathbb{C} \setminus \{0, 1, t\}$$

whose fibers are the affine curves Γ_s , (5), $s \in \mathbb{C} \setminus \{0, 1, t\}$. Each Γ_s is topologically a torus with two removed points. Hence $\dim H_1(\Gamma_s, \mathbb{Z}) = \dim H_{DR}^1(\Gamma_s, \mathbb{C}) = 3$. Therefore if $\gamma(s) \in H_1(\Gamma_s, \mathbb{Z})$ is a family of cycles depending continuously on s , then the Abelian integral

$$I(s) = \int_{\gamma(s)} \omega, \omega = P(\xi, \eta)d\xi + Q(\xi, \eta)d\eta, P, Q \in \mathbb{C}[\xi, \eta]$$

satisfies a Fuchsian differential equation of order three, whose coefficients are polynomials in s, a . In the case when the differential form ω has no residues, it satisfies a second order equation. Explicitly, if $\gamma_1(s), \gamma_2(s)$, is a continuous family of cycles generating the homology group of the compactified elliptic curve Γ_s , then the equation reads

$$\det \begin{pmatrix} x & x' & x'' \\ \int_{\gamma_1(s)} \omega & (\int_{\gamma_1(s)} \omega)' & (\int_{\gamma_1(s)} \omega)'' \\ \int_{\gamma_2(s)} \omega & (\int_{\gamma_2(s)} \omega)' & (\int_{\gamma_2(s)} \omega)'' \end{pmatrix} = 0.$$

It follows from the Picard-Lefschetz formula and the moderate growth of the integrals, that the coefficients of the above differential equations are rational in s, a . A local analysis of the singularities shows for instance that

$$\det \begin{pmatrix} \int_{\gamma_1(s)} \omega & (\int_{\gamma_1(s)} \omega)' \\ \int_{\gamma_2(s)} \omega & (\int_{\gamma_2(s)} \omega)' \end{pmatrix} = \frac{p(s, a)}{s(s-1)(s-t)}$$

where $p(s, a)$ is a polynomial in s, a . If we put $\omega = dx/y$ then $\int_{\gamma_1(s)} \omega$ grows no faster than $s^{1/4-1/2}$ at ∞ (for a fixed a). Thus

$$\frac{p(s, a)}{s(s-1)(s-t)}$$

grows at infinity no faster than $s^{-1/2-1}$ and hence no faster than s^{-2} . It is expected therefore that $p(s, a)$ is of degree one in s and the corresponding root, which we denote by λ , is an apparent singularity for the Picard-Fuchs equation in consideration. We are therefore in a position to apply Theorem 2, provided that the deformation of the Fuchs equation with respect to

the parameter a is isomonodromical. Indeed, the monodromy group of our equation is contained in $SL(2, \mathbb{Z})$ which shows that any deformation of this equation is isomonodromical. The Picard-Lefschetz formula shows that the monodromy group in question is generated, up to conjugacy, by the matrices

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}. \quad (7)$$

To deduce an explicit formula for the corresponding algebraic solution of \mathcal{P}_{VI} we need explicit formulae for the Picard-Fuchs equations.

Lemma 1 *Let $\gamma(s) \in H_1(\Gamma_s, \mathbb{Z})$ be a family of cycles depending continuously on s . The complete elliptic integrals of first and second kind*

$$x(s) = \int_{\gamma(s)} \frac{d\xi}{\eta}, \quad y(s) = \int_{\gamma(s)} \frac{(3\xi^2 - 2(a+1))\xi d\xi}{\eta}$$

satisfy Picard-Fuchs equations of the form

$$a_0(s)x'' + a_1(s)x' + a_2(s)x = 0$$

$$b_0(s)y'' + b_1(s)y' + b_2(s)y = 0$$

where

$$\begin{aligned} a_0(s) &= s(s-1)((2a-1)s + a^3(a-2))((a^2-a+1)s + a^2(a-2)) \\ a_1(s) &= 2(2a-1)(a^2-a+1)s^3 + (a^6-3a^5+9a^4-19a^3+9a^2-3a+1)s^2 \\ &\quad + 2a^2(a-2)(a^4-2a^3-2a+1)s - a^5(a-2)^2 \\ a_2(s) &= (2a-1)[27(a^2-a+1)s^2 - (a-2)(2a^4-a^3-60a^2-a+2)s \\ &\quad + a^2(a-2)^2(10a^2+11a+10)]/144 \\ b_0(s) &= s(s-1)((2a-1)s + a^3(a-2))((a^2-7a+1)s - a(a-2)(2a^2+a+2)) \\ b_1(s) &= (2a-1)s[(a^2-7a+1)s^2 - 2a(a-2)(2a^2+a+2)s \\ &\quad - a(a-2)^2(a^4+a^3+a^2+a+1)] \\ b_2(s) &= -(2a-1)[9(a^2-7a+1)s^2 - (a-2)(10a^4+31a^3-12a^2+31a+10)s \\ &\quad - a(a-2)^2(2a^2+a+2)]/144 \end{aligned}$$

The proof of the above Lemma is straightforward, see for instance [5]. It is seen that the roots of $a_0(s)$ are 0, 1 and

$$\lambda = \frac{a^2(2-a)}{a^2-a+1}, \quad t = \frac{a^3(2-a)}{2a-1}$$

which implies the algebraic solution (3). In the same way the roots of $b_0(s)$ provide the solution (4). The Riemann schemes of the Picard-Fuchs equations for $x(s), y(s)$ are given by

$$\begin{pmatrix} 0 & 1 & \frac{a^3(2-a)}{2a-1} & \frac{a^2(2-a)}{a^2-a+1} & \infty \\ 0 & 0 & 0 & 0 & \frac{1}{4} \\ 0 & 0 & 0 & 2 & \frac{3}{4} \end{pmatrix}$$

and

$$\begin{pmatrix} 0 & 1 & \frac{a^3(2-a)}{2a-1} & \frac{a(a-2)(2a^2+a+2)}{a^2-7a+1} & \infty \\ 0 & 0 & 0 & 0 & \frac{1}{4} \\ 1 & 0 & 0 & 2 & -\frac{1}{4} \end{pmatrix}.$$

The Corollary after Theorem 2 implies that the curve (3) is an integral curve of the $\mathcal{P}_{VI}(\alpha)$ equation with parameters $(\alpha_0, \alpha_1, \alpha_2, \alpha_3) = (\frac{1}{8}, 0, 0, 0)$, see (6). Similarly, the Fuchsian equation satisfied by the complete elliptic integral of second kind $y(s)$ provides the algebraic solution (4) of $\mathcal{P}_{VI}(\alpha)$ with $\alpha = (\frac{1}{8}, \frac{1}{2}, 0, 0)$.

It is remarkable that (3) was found to be a solution of $\mathcal{P}_{VI}(\alpha)$ by Hitchin [6][p.177], but for $\alpha = (\frac{1}{8}, \frac{1}{8}, \frac{1}{8}, \frac{1}{8})$. After taking the difference between these two equations we obtain the following affine equation of the integral curve (3)

$$-\frac{t}{\lambda^2} + \frac{t-1}{(\lambda-1)^2} - \frac{t(t-1)}{(\lambda-t)^2} = 0.$$

This also shows that (3) is a common algebraic solution of the pencil of $\mathcal{P}_{VI}(\alpha)$ equations

$$\alpha = (\frac{1}{8}, \frac{s}{8}, \frac{s}{8}, \frac{s}{8}), s \in \mathbb{C}.$$

This completes the proof of Theorem 1.

References

- [1] V. ARNOLD, A. VARCHENKO, S. GOUSSEIN-ZADÉ, *Singularités des applications différentiables, 2^{eme} partie*, Edition Mir, Moscou, 1986
- [2] B. Ben Hamed, *Connexion de Gauss-Manin et Solutions Algébriques de \mathcal{P}_{VI}* , mémoire de DEA, Université de Toulouse III, 2003.

- [3] Ph. Boalch, The Klein solution to Painlevé's sixth equation, math.AG/0308221 v2 (2003).
- [4] B. Dubrovin, M. Mazzocco, Monodromy of certain Painlevé-VI transcendents and reflection groups, Invent. Math **141**, 55-147 (2000).
- [5] F. DUMORTIER, CH. LI, Perturbations from an elliptic Hamiltonian of degree four, J. Diff. Equations, **176**, 114-157 (2001).
- [6] N. J. HITCHIN, Poncelet polygons and the Painlevé equations, in *Geometry and Analysis*, (S. Ramanan, ed.) Oxford University Press, Bombay, 1995.
- [7] K.IWAZAKI, H.KIMURA, S.SHIMOMURA, M.YOSHIDA, *From Gauss to Painlevé*, Aspects of Mathematics, Vieweg 1991.
- [8] YU. I. MANIN, *Sixth painlevé equation, Universal elliptic curve and Mirror of \mathbb{P}^2* , Amer. Math. Soc. Transl. (2) Vol. **186**, 1998.